AUG 18 2003 00:49 FR PENNIE & EDMONDS

212 869 9741 TO 17037465060

P.76 #}

eceived attent

5/18/03

DE

EXIDELL

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Michael Meyrick Burrell and

Stephen Andrew Coates

Serial No.: 08/192,493

E & EDMONDS

5,830,724

Filed:

February 7, 1994

Art Unit:

.1804

For:

MODIFICATION OF STARCH PRODUCTION

Examiner:


Charles C.P. Rories


DECLARATION UNDER RULE 37 CFR 1.132

- I, Michael Meyrick BURRELL, of 20 Histon Road, Cottenham, Cambridge, England, DO SOLEMNLY and SINCERELY DECLARE as follows:
- 1. I am the Michael Meyrick Burrell who is named as being an Applicant in respect of United States Patent Application Serial No. 08/192,493.
- 2. I am a scholar of Cambridge University and hold a PhD in Plant Biochemistry and a Degree in Natural Sciences with honours in Botany.
- I initiated and supervised the conduct of the experiment now to be related.

E & EDMONDS

- 2
- 4. The coding sequence for the wheat homologue of waxy was fused to the high molecular weight glutenin promoter of wheat to produce a chimaeric gene which would be expressed in developing endosperm. A hybrid maize was transformed with this construct using the particle bombardment method and plants were regenerated to produce seed by outcrossing.
- 5. Analysis of single seeds taken from six independent transformants is shown below. Starch was isolated and the starch granule bound proteins separated by gel electrophoresis. The 60kDa waxy protein is clearly absent from line 3 (Figure 1). The starch from this line clearly has a much reduced amylose content (Figure 2).





Received from < 212 869 9741 > at 8/18/03 5:04:16 PM [Eastern Daylight Time]

۵. ہد. ب

LE & EDMONDS

The seed from the progeny of plants obtained in the 6. transformation procedure were divided into two populations. One population contained plants that showed no clear 60kDa waxy protein product and one population contained segregants that had the 60kDa waxy protein. A third population of untransformed controls were also included in the analysis. results (table below) clearly show the somewhat surprising result that the absence of the waxy protein is associated with no detectable amylose and no change in starch content.

	Starch Content umol/gfwt		% amylose		
Grain analysed Control line	Mean	S.E.M.	Mean	S.E.M.	Number of grain analysed
grain	91.30593	13.83537	19.28946	1.834417	5
Progeny grain lacking the product of the waxy gene	105.9155	11.64986	nd*	nd*	15
Progeny grain expressing the product of the waxy gene	92.26848	8.171925	21.92222	0.71156	16

*nd == not detected

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that wilful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code,

E & EDMONDS 212 869 9741 TO 179

and that such wilful false statements may jeopardise the validity of the application or any patent issuing thereon.